



Lip Liner and Full Lip Color Medication

If you have ever had a cold sore or fever blister in the past, you harbor the herpes simplex virus within your body. Every person is different in his or her activity of this virus. Some may have frequent outbreaks, and some may have only had one or two outbreaks a long time ago, nevertheless, the virus is still harbored within your body. The lip liner and full lip color procedure may cause an outbreak in some individuals who have this virus. There fore you are advised to acquire the below mentioned medication, prior to your procedure, to alleviate the possibility of an outbreak. This is an optional medication and Contemporary Plastic Surgery does not hold any liability regarding its use or nonuse for prevention of cold sores/fever blisters.

*Please request **one** of these anti-viral medications from your Physician:*

Zovirax (Acyclovir)

Famvir

Valtrex

*Your will need to take the prescribed dosage of **one** of the medication up above for **five** (5) consecutive days. Begin by taking the pills **two** (2) **days before** your lip procedure, the **same day** as your procedure and **two** (2) **days following** your lip procedure.*

Remember: *Each of these medications has a different dosage, so please check with your physician regarding adequate dosing and administration.*

Please request enough medication to cover you for two (2) lip procedures.

*I, _____, have been advised to take this medication for prophylaxis purposes due to my susceptibility to the herpes virus. By taking this medication for a lip liner or full lip color procedure, I am aware that it does not guarantee that I will not have a herpes outbreak, and do not hold **Contemporary Plastic Surgery** responsible if this should occur.*

Signature

Date